



The Charge Form Factor of the Neutron at Low Q^2

- Introduction/Motivation
- BLAST Experiment
- Results and Discussion



The Charge Form Factor of the Neutron at Low Q^2





Experimental Program

High quality data for nucleon and deuteron structure
by means of *spin-dependent electron scattering*

Pol. H	$\vec{p}(\vec{e}, e')$ Gen. GDH	$\vec{p}(\vec{e}, e'p)$ G^p_E/G^p_M	$\vec{p}(\vec{e}, e'\pi^+)n, \vec{p}(\vec{e}, e'p)\pi^0, \vec{p}(\vec{e}, e'n)\pi^+$ N- Δ : C2/M1	
Vect-Pol. D	$\vec{d}(\vec{e}, e')$ G^n_M	$\vec{d}(\vec{e}, e'p)$ D-state	<div>$\vec{d}(\vec{e}, e'n)$ G^n_E</div>	$\vec{d}(\vec{e}, e'd)$ T^e_{11}
Tens-Pol. D	$\vec{d}(e, e'd)$ T_{20}	$\vec{d}(e, e'p)$ D-state		




Scientific Motivation

- Nucleon form factors provide basic information on nucleon structure
- G_E^n is the least known among the nucleon form factors, with errors of typically 15-20%
- Low- Q^2 region is a testing ground for QCD and pion-cloud inspired and other effective nucleon models
- G_E^n related to neutron charge distribution
- Precise knowledge of G_E^n is essential for parity violation experiments




MIT-Bates Linear Accelerator Center




- **Linac:** 2×500 MeV
- **Beam:** 850 MeV / $I_{\max} = 225$ mA / $P_e = 65$ %
- **SHR:** Siberian Snake + Compton Polarimeter
- **Target:** Internal Target = Atomic Beam Source



Atomic Beam Source






- Isotopically pure H or D
 - Vector Polarized H
 - Vector and Tensor D
- Target Thickness/Luminosity
 - Flow 2.2×10^{16} atoms/s
 - Density 6×10^{13} atoms/cm²
 - Luminosity 6×10^{31} cm⁻²s⁻¹
- Target Polarization typically 70-80%






Atomic Beam Source






Atomic Beam Source




- Quasielastic $^2\vec{H}(\vec{e}, e'n)p$
- Beam-Target Asymmetry
- $A_{\text{ed}}^{\text{v}}(\text{exp}) = h \cdot P_z \cdot A_{\text{ed}}^{\text{v}}(\text{th})$
- $\langle h P_z \rangle = 0.567 \pm 0.006$
 $\langle P_z \rangle = 0.85 \pm 0.04$
 $\langle h \rangle = 0.67 \pm 0.04$






BLAST Detectors

- Left-Right symmetry
- Large Acceptance
 - $0.1 < Q^2/(\text{GeV}/c)^2 < 1.0$
- Coils: $B = 3.8 \text{ kG}$
- Drift Chambers
 - PID, tracking
 - $\delta\theta \approx 0.5^\circ$, $\frac{\delta p}{p} \approx 3\%$
- Cerenkov Counters
 - e, π separation
- Scintillators
 - TOF, PID, trigger
- Neutron Counters
 - Neutron ToF





Kinematics and Observables




- Electrodissintegration of the Deuteron
- Quasi-elastic $^2\text{H}(e,e'n)$
- Beam + Target Polarized

$$\frac{d\sigma}{d\Omega_e dE_{e'} d\Omega_{CM}} = S_0 (1 + P_d^V A_d^V + P_d^T A_d^T + h(A_e + P_d^V A_{ed}^V + P_d^T A_{ed}^T))$$

$$A_{ed}^V = \frac{a G_M^{n^2} \cos \theta^* + b G_E^n G_M^n \sin \theta^* \cos \phi^*}{c G_E^{n^2} + G_M^{n^2}} \approx a \cos \theta^* + b \frac{G_E^n}{G_M^n} \sin \theta^* \cos \phi^*$$




Experimental Layout





Experimental Layout



Electron - Right Sector


“Parallel” Kinematics



$$\theta^* \approx 0$$



Experimental Layout



Electron - left sector

“Perpendicular” Kinematics



$$\theta^* \approx 90$$




Blast data

- 3 MC integrated charge delivered to BLAST
- Programs for polarized hydrogen and vector/tensor polarized deuterium
- Deuterium run May-October 2004, spin angle 32°
450 kC charge (169 pb^{-1}), $P_z=85\%$, $P_{zz}=66\%$
- Deuterium run March-May 2005, spin angle 47°
550 kC charge (150 pb^{-1}), $P_z=70\%$, $P_{zz}=54\%$
- Preliminary data will be presented for 2004 run



Invariant Mass and Time of Flight


- Very clean quasi-elastic ${}^2\text{H}(e,e'n)p$ spectrum
- Highly efficient proton veto (Wire Chambers)





Invariant Mass and Time of Flight


- Very clean quasi-elastic ${}^2\text{H}(e,e'n)p$ spectrum
- Highly efficient proton veto (Wire Chambers)





Extraction of G_E^n


- Quasielastic $^2\vec{H}(\vec{e}, e'n)p$
- Full Monte Carlo Simulation of the BLAST experiment
- Deuteron Electrodisintegration cross section calculations by **H. Arenhövel**
- Accounted for FSI, MEC, IC, RC
- Spin-perpendicular beam-target asymmetry $A_{ed}^V(90^\circ, 0^\circ)$ shows high sensitivity to G_E^n





Extraction of G_E^n

- Quasielastic $^2\vec{H}(\vec{e}, e' n)p$
- Full Monte Carlo Simulation of the BLAST experiment
- Deuteron Electrodisintegration cross section calculations by **H. Arenhövel**
- Accounted for **FSI, MEC, IC, RC**
- Spin-perpendicular beam-target asymmetry $A_{ed}^V(90^\circ, 0^\circ)$ shows high sensitivity to G_E^n
- Compare measured A_{ed}^V with **BLASTMC**, varying G_E^n






Systematic Errors

<hr/>	
■ Uncertainty of target spin angle	5%
□ 12% per degree	
■ Beam-target polarization product	2.5%
■ Radiative effects	<1.0%
□ Small helicity dependency	
■ Uncertainty of G_M^{η}	1.5%
■ Model dependency	<3%
□ Effect of potential negligible	
□ Final state interaction reliable	
Total:	6.6%



Results and Discussion


- World data on G_n^E from double pol. Experiments






Results and Discussion

- World data on G_n^E from double pol. Experiments
- Including BLAST 2004





Results and Discussion




- World data on G_E^n from double pol. Experiments
- Including BLAST 2004
- BLAST Fit

□ $\langle r_n^2 \rangle = -0.115 \text{ fm}^2$




Nucleon Models



- World data on G_E^n from double pol. Experiments




Nucleon Models



- World data on G_E^n from double pol. Experiments
- **Dispersion Theory**




Nucleon Models



- World data on G_E^n from double pol. Experiments
- Dispersion Theory
- Chiral Soliton




Nucleon Models



- World data on G_E^n from double pol. Experiments
- Dispersion Theory
- Chiral Soliton
- R-CQM




Nucleon Models



- World data on G_E^n from double pol. Experiments
- Dispersion Theory
- Chiral Soliton
- R-CQM
- Cloudy bag




Nucleon Models



- World data on G_E^n from double pol. Experiments
- **Dispersion Theory**
- **Chiral Soliton**
- **R-CQM**
- **Cloudy bag**
- **Diquark**



Nucleon Models



- World data on G_E^n from double pol. Experiments
- Dispersion Theory
- Chiral Soliton
- R-CQM
- Cloudy bag
- Diquark
- Lattice



Conclusion

- Measure G_E^n with quasielastic scattering of polarized electrons from vector-polarized deuterium using BLAST at MIT-Bates
- Very small systematic errors
- G_E^n overall known to $\approx 5\%$ at $Q^2 < 1 \text{ (GeV/c)}^2$
- Dispersion theory gives the best description
- No theory describes G_E^n at low and high Q^2 simultaneously
- Evidence for enhancement at low Q^2 - role of pion cloud?



Outlook

- Only **half** of the BLAST data analyzed so far, prel. results of full dataset envisioned for **December 2005**
- With new precision data of T_{20} from BLAST and with **improved $A(Q^2)$** new attempt to G_E^n determine from G_Q
- **ed elastic analysis:** Mainz-Saclay discrepancy **8% in $A \Rightarrow$ factor of 2 in G_E^n**
 - New measurements of $A(Q^2)$ at JLab (E-05-004) and MAMI
- Extension of G_E^n at high $Q^2 < 3.5 \text{ (GeV/c)}^2$ (E-02-13)
- Proposal of **BLAST@ELSA/Bonn**
Measure G_E^n for $Q^2 = 0.04\text{-}1.5 \text{ (GeV/c)}^2$ with both vector-polarized ^2H and polarized ^3He